

MEMOIRS OF THE QUEENSLAND MUSEUM

BRISBANE

© Queensland Museum
PO Box 3300, South Brisbane 4101, Australia
Phone 06 7 3840 7555
Fax 06 7 3846 1226
Email qmlib@qm.qld.gov.au
Website www.qm.qld.gov.au

National Library of Australia card number
ISSN 0079-8835

NOTE

Papers published in this volume and in all previous volumes of the *Memoirs of the Queensland Museum* may be reproduced for scientific research, individual study or other educational purposes. Properly acknowledged quotations may be made but queries regarding the republication of any papers should be addressed to the Editor in Chief. Copies of the journal can be purchased from the Queensland Museum Shop.

A Guide to Authors is displayed at the Queensland Museum web site

A Queensland Government Project
Typeset at the Queensland Museum

SOUTHERLY RANGE EXTENSION FOR COMMON ROCK-RAT *ZYZOMYS ARGURUS* (RODENTIA: MURIDAE) IN QUEENSLAND. *Memoirs of the Queensland Museum* 48(1): 154. 2002.- The Common Rock-rat (*Zyzyomys argurus* Thomas, 1889) is a small to medium-sized rodent upto 55g; it is almost always associated with rocky outcrops (Fleming, 1995); it is endemic to N Australia (WA, NT, QLD); in QLD it occurs in rocky ranges and outcrops between Lagoon Creek Gorge, (17°33'S, 138°01'E) and Cape Melville, (14°22'S, 144°37'E) to Lark Quarry, (23°01'S, 142°24'E) and Blackdown Tableland, (23°44'40"S, 149°06'20"E) (Kitchener, 1989; QM data).

In May 2000 a trapping survey for Northern Quoll (Oakwood & Firestone, 2000), in the Mount Moffatt Section of Carnarvon NP, 600km NW of Brisbane used wire cage, and Type A and Type B Elliott traps set at 7 sites (454 trap nights) on or near rocky outcrops. Traps were baited with a bolus of rolled oats and peanut butter, or salami laced with pistachio essence. *D. hallucatus*, Fawn-footed Melomys *Melomys cervinipes*, Long-nosed Bandicoot *Perameles nasuta*, *Pseudomys* sp. and *Z. argurus* were caught.

This new record for *Z. argurus* was caught at The Duchess (25°06'11.6"S, 147°51'06.6"E), a distinctive outcrop of Precipice Sandstone overlooking the Maranoa River and rising above the surrounding sandplain to an altitude of 740m. Two *D. hallucatus* and one unidentified rodent were also caught at this site (116 trap nights). *Z. argurus* was captured on 30 May 2000 in a collapsible wire cage trap (55.5x20x20cm), baited with peanut butter and rolled oats. The trap was set at the entrance of a cave in the vertical rock-face, on a broad ledge just below the summit. The cave faces SW, is up to 2.0m high, 1.55m wide and 13.8m long. No evidence of *Z. argurus*, such as scats or chewed seeds and fruit, was observed in the vicinity of the trap site.

Vegetation in the trap site is scattered *Angophora leiocarpa* and *Eucalyptus decorticans*, with a mid-layer dominated by *Leptospermum lamellatum*, *Acacia* spp., *Callitris endlicheri* and low shrubs. The ground layer is dominated by *Lomandra* sp., while other graminoids and forbs are sparse. Dense leaf litter covers the ground. Areas of vertical rock-face are dominated by *L. lamellatum*, *Wahlenbergia islensis*, and bryophytes. The area equates to regional ecosystem 11.10.4 'complex of *Eucalyptus decorticans* and/or *Acacia shirleyi* woodland/open forest and mixed woodland/open forest on Cainozoic to Proterozoic consolidated, medium to coarse grained sediments' (Young et al., 1999). This habitat type is extensive in Mount Moffatt, Moolayember, Salvator Rosa and Goodliffe Sections of Carnarvon National Park, and is broadly consistent with habitat records for *Z. argurus* in other parts of Queensland.

Z. argurus was extremely active in the trap when first approached and the tip of the tail was discarded when the animal was first handled. The voucher specimen, lodged in the Queensland Museum (QMJM 14298), is a juvenile male, with SVL ~85mm and tail length ~87mm. Its body has brownish-grey fur above, buff on the sides and cream below. The tail is distinctly bicoloured (dark grey above, light grey below) and slightly, but noticeably swollen near the base, clearly ringed (as opposed to mosaic) and well furred. The nose is pink, ears dark grey and feet dark grey, furred white above. The specimen was distinguished from the other 8 eight species of rodent that occur at Mount Moffatt by a

combination of body size, SVL/tail length ratio, un-notched upper incisors, strongly bi-coloured well furred tail which is swollen at the base, and pronounced roman-nose.

The southern limit for *Z. argurus* was Blackdown Tableland (23°44'40"S, 149°06'20"E) where it was trapped between Peregrine and Horseshoe Lookouts, Sept.1984 (C. James, G. Porter pers. comm.). This record in the Carnarvon Range represents a range extension of 200km. The climate of Carnarvon Range is subhumid, with mean annual rainfall of >800mm (Galloway et al., 1974). The vertebrate fauna is predominantly Bassian whereas *Z. argurus* is one of a small subset (7 spp.) of Torresian species.

Zyzyomys argurus is regarded as secure at a national level (Lee, 1995), and common in Queensland (Dickman et al., 2000).

Acknowledgements

Thanks to Rick McKiernan, Liam Oakwood, Warwick Greville, Karen Firestone and Toby Ware for field assistance. Greg Mifsud, Gary Porter and Cameron James provided unpublished *Z. argurus* records. Richard Johnson, Rod Hobson, Wayne Houston and Dr. David McFarland commented on early drafts. This research was conducted under Queensland Parks and Wildlife Service permit W0/002493/00/SAA (to M. Oakwood).

Literature Cited

- DICKMAN, C.R., LEUNG, L. K.-P & VAN DYCK, S.M. 2000. Status, ecological attributes and conservation of native rodents in Queensland. *Wildlife Research* 27: 333-346.
- FLEMING, M. 1995. Common Rock-rat *Zyzyomys argurus*. Pp. 620-621. In Strahan, R. (ed) *The Mammals of Australia*. (Australian Museum/Reed: Sydney).
- GALLOWAY, R.W., GUNN, R.H., PEDLEY, L., COCKS, K.D. & KALMA, J.D. 1974. Lands of the Balonne-Maranoa area, Queensland. Land Research Series No. 34. (CSIRO: Melbourne).
- KITCHENER, D.J. 1989. Taxonomic appraisal of *Zyzyomys* (Rodentia, Muridae) with descriptions of two new species from the Northern Territory, Australia. *Records of the Western Australian Museum* 14: 331-373.
- LEE, A.K. 1995. The Action Plan for Australian Rodents. (Australian Nature Conservation Agency: Canberra).
- OAKWOOD, M. & FIRESTONE, K. 2000. Conservation and Management of the Northern Quoll *Dasyurus hallucatus*. Unpublished report to the Queensland Parks and Wildlife Service.
- YOUNG, P.A.R., WILSON, B.A., McCOSKER, J.C., FENSHAM, R.J., MORGAN, G. & TAYLOR, P.M. 1999. Brigalow Belt. Chapter 11. In Sattler, P.S. & Williams, R.D. (eds) *The Conservation Status of Queensland's Bioregional Ecosystems*. (Environmental Protection Agency: Brisbane).

Craig Eddie, Environmental Protection Agency, PO Box 981, Roma 4455, Australia; M. Oakwood, Department of Ecosystem Management, University of New England 2351, Australia; 10 August 2001.